



431 E. Locust Street Des Moines, IA 50309 Phone (515) 309-4808 Fax (515) 243-2880

PHYSICIAN OR OTHER AUTHORIZED HEALTH PROFESSIONAL ORDER FORM FOR THERAPEUTIC PHEBOTOMY

Physician or Other Authorized Health Professional Section

Diagnosis: Hereditary Hemochromatosis, Polycythemia, Polycythemia Secondary to Testosterone Therapy, Other. Phlebotomy Frequency: Every \_\_\_ week(s) for \_\_\_ weeks, Every \_\_\_ month(s) for \_\_\_ months, One time only, Other: Specify frequency and duration. Amount to Draw at Each Phlebotomy: 500 mls will be drawn unless specified below. DELAY OR CANCEL PHEBOTOMY IF HEMOGLOBIN IS BELOW: Hgb value. Please Note: \*LifeServe Blood Center does NOT perform Ferritin testing. Physician or Other Authorized Health Professional Signature: Date:

STANDING ORDERS MUST BE RENEWED ANNUALLY

PRINT PHYSICIAN or OTHER AUTHORIZED HEALTH PROFESSIONAL INFORMATION

Physician or Other Authorized Health Professional Name

Physician or Other Authorized Health Professional Address STREET CITY STATE ZIP

Physician or Other Authorized Health Professional Phone # FAX#

PRINT PATIENT INFORMATION

Patient Name: LAST FIRST MI

DOB: Patient Phone #

LIFESERVE BLOOD CENTER USE ONLY

Medical Director Approval not required ID# Date

Donor ID# Order Expiration Date: APPROVAL MEDICAL DIRECTOR DATE:

INFORMATION FOR THE ORDERING PHYSICIAN OR OTHER AUTHORIZED HEALTH PROFESSIONAL

There is a 2-3% incidence of vasovagal reactions with the donation of one unit. A severe vasovagal reaction with marked hypotension and syncope may compromise a patient's cardiac or cerebrovascular status if occlusive disease is already present. There are rare reports of myocardial infarctions following donations, even in "healthy" blood donors. The benefits of the therapeutic phlebotomy should outweigh the risks. The requesting physician or other authorized health professional should determine the safety of donation by the patient. Donation associated with a severe vasovagal reaction during pregnancy could result in decreased blood flow to the fetus. These types of problems must be addressed in the written order, or, if the problem is unfamiliar to the ordering physician, a note of approval for phlebotomy from a physician familiar with the problem will suffice. The Medical Director at LifeServe Blood Center may be contacted for advice on the safety of donation for certain individuals. The patient's health status must be medically stable enough to tolerate the therapeutic phlebotomy process. The patient's blood pressure, pulse and hemoglobin/hematocrit are checked prior to each phlebotomy. A report will be sent to the ordering physician or other authorized health professional. The patient MUST call to make an appointment. This service is offered on specific days and specific times, depending on location. If you have questions regarding the Therapeutic Phlebotomy Program, call (515) 309-4808.